

279-3380

10/02/2012

1/29



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 2, 2012

Timothy M Formella
Product Registration Manager
FMC Corporation
1735 Market Street
Philadelphia, PA 19103

Subject Amendment – Removing conflicting application restriction on corn
F9043 EC Insecticide
EPA Reg No 279-3380
Your submission dated July 20, 2012

Dear Mr Formella

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable. A stamped copy of the label is enclosed for your records.

If you have any questions regarding this action, please contact BeWanda Alexander at Alexander.bewanda@epa.gov or (703) 305-7460

Sincerely,

A handwritten signature in black ink, which appears to read "BeWanda Alexander" but is likely the signature of Richard Gebken, as indicated by the typed name below.

Richard Gebken
Product Manager Team 10
Insecticide Branch
Registration Division (7505P)

Enclosure

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms
For retail sale to and use only by certified applicators or
persons under their direct supervision and only for those uses
covered by the certified applicator's certificate

F9043 EC Insecticide**EPA Reg No 279-3380****EPA Est 279-****Active Ingredients****By Wt**

Zeta Cypermethrin*

8 20%

Bifenthrin**

9 80%

Other Ingredients***82 00%


100 0%

F9043 EC Insecticide contains 1.5 pounds active ingredients per gallon

* *Cis/trans* isomer ratio Max 75% (\pm) *cis* and Min 25% (\pm) *trans** *Cis* isomers 97% minimum *trans* isomers 3% maximum

* * Contains Petroleum Distillates

U S Patent No Pending

**KEEP OUT OF REACH OF CHILDREN****WARNING
AVISO**

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label
find someone to explain it to you in detail)

See other panels for additional precautionary information

FMC

FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia PA 19103

NET CONTENTS

FIRST AID	
IF SWALLOWED	Immediately call a poison control center or doctor Do not induce vomiting unless told to do so by a poison control center or doctor Do not give any liquid to the person Do not give anything by mouth to an unconscious person
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eyes Call a poison control center for treatment advice
IF ON SKIN OR CLOTHING	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 to 20 minutes Call a poison control center or doctor for treatment advice
IF INHALED	Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible Call a poison control center or doctor for further treatment advice
NOTE TO PHYSICIAN	
Contains petroleum distillates Vomiting may cause aspiration pneumonia	
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 (800) 331 3148 for Emergency Assistance	

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning

May be fatal if swallowed Harmful if inhaled or absorbed through the skin Causes moderate eye irritation Avoid contact with skin eyes or clothing Avoid breathing spray mist Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or before using the toilet Remove and wash contaminated clothing before reuse

Personal Protective Equipment

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for category G on an EPA chemical resistance category selection chart

Handlers who may be exposed to the dilute through application or other tasks must wear Long sleeved shirt and long pants chemical resistant gloves such as Barrier Laminate or Butyl Nitrile or Neoprene Rubber ≥ 14 mls or Viton ≥ 14 mls and shoes plus socks

Handlers who may be exposed to the concentrate through mixing loading application or other tasks must wear Long sleeved shirt and long pants chemical resistant gloves such as Barrier Laminate or Butyl Nitrile or Neoprene Rubber ≥ 14 mls or Viton ≥ 14 mls and shoes plus socks and protective eyewear

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate Do not reuse them

User Safety Recommendations

Wash thoroughly with soap and water after handling. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (540CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

This pesticide is extremely toxic to fish, aquatic invertebrates, oysters, and shrimp. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

The use of F9043 EC Insecticide is prohibited in areas where its application may result in exposure to endangered species. Prior to use in a particular county, contact the local extension service for procedures and precautions to use to protect endangered species.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

Physical or Chemical Hazards

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Resistance. Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves such as barrier laminate or Viton, and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water food or feed or other pesticides or fertilizers by storage and disposal

Pesticide Storage

Do not freeze Keep out of reach of children and animals Store in original containers only Store in a cool dry place and avoid excess heat Carefully open containers After partial use replace lids and close tightly Do not put concentrate or dilute material into food or drink containers

In case of spill avoid contact isolate area and keep out animals and unprotected persons Confine spills Call CHEMTREC (Transportation and Spills) (800) 424 9300

To confine spill Dike surrounding area or absorb with sand cat litter or commercial clay Place damaged package in a holding container Identify contents

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

Container Handling

Nonrefillable Container

Do not reuse or refill this container

Triple rinse container (or equivalent) promptly after emptying

Triple rinse as follows

For containers equal to 5 gallons or less Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal

Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times

For containers greater than 5 gallons Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times

Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or incineration or if allowed by state and local authorities by burning If burned stay out of smoke

Refillable/Returnable Container

Refill this container with pesticide only Do not reuse this container for any other purpose

Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean the container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10% full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times

Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or incineration or if allowed by state and local authorities by burning If burned stay out of smoke

Chemigation Use Directions

Apply this product only through sprinkler including center pivot lateral move end tow side (wheel) roll traveler big gun solid set or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from non uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must also contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

F9043 EC Insecticide should be applied continuously for the duration of the water application. F9043 EC Insecticide should be diluted in sufficient volume to insure accurate application over the area to be treated.

Use the appropriate amount of water to carry the product to the target pest. Agitation is not required when a suitable diluent is used.

BUFFER ZONES

Vegetative Buffer Zones

Construct and maintain a minimum 10 foot wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, estuaries, and commercial fish farm ponds).

Only apply products containing zeta cypermethrin and bifenthrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses, Natural Resources Conservation Services, USDA NRCS, 2000 Fort Worth, Texas, 21pp. <http://www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf>

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast) – Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for ULV Aerial Application – Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for Non ULV Aerial Application – Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Spray Drift Requirements

Wind Direction and Speed

Only apply this product if the wind direction favors on target deposition. Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

Do not make aerial or ground applications into temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only Medium or coarser spray nozzles (for ground and non ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Preventive Use

For cutworm, armyworm, grape colaspis, or stalk borer control, F9043 EC Insecticide may be applied before, during, or after planting. For soil incorporated applications, use higher rates for improved control.

Adjuvants

Adjuvants may be used with F9043 EC Insecticide. When selecting an adjuvant, select an adjuvant that meets or exceeds CPDA Adjuvant Certification. It is recommended for optimum performance. Refer to the individual crop recommendation sections of this label for specific adjuvant type and use rates.

Rotational Crops

Crops for which bifenthrin and zeta cypermethrin tolerances exist may be rotated at any time. All other crops may be rotated 30 days following the final application of F9043 EC Insecticide.

Tank-Mixture

F9043 EC Insecticide may be applied in tank mixtures with other products approved for use on the F9043 EC Insecticide and tank mix partners' labels. Observe all restrictions and precautions that appear on the labels of these products. Test for compatibility of products before mixing.

F9043 EC Insecticide contains the pyrethroids zeta-cypermethrin and bifenthrin

Maximum Usage When Applying Zeta Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season

Do not apply more than the maximum seasonal total for either zeta cypermethrin or cypermethrin products when used alone. Do not apply more than the combined maximum seasonal total for both products as outlined in the table below.

Maximum Seasonal Total

Maximum Usage When Applying Zeta Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season

Crop	Maximum Seasonal Total (lbs ai/acre)			Maximum Seasonal Total (lbs ai/acre) When Applying Cypermethrin and Zeta cypermethrin Products to the Same Crop		Maximum Seasonal Total (lbs ai/acre) When Applying Zeta-cypermethrin Products to the Same Crop
	Zeta cypermethrin			Cypermethrin	Zeta cypermethrin plus Cypermethrin	Zeta cypermethrin
	Mustang	Mustang Max	F9043			
Canola	0.3	0.15	0.07	NA	NA	0.3
Field Corn	0.2	0.10	0.10	NA	NA	0.2
Sweet Corn	0.3	0.15	0.15	NA	NA	0.3
Cotton	0.3	0.15	0.15	0.6	0.6	0.3
Peanut	0.3	0.15	0.15	NA	NA	0.3
Potato	0.3	0.15	0.15	NA	NA	0.3
Soybean	0.3	0.15	0.15	NA	NA	0.3
Head and Stem Brassica	0.3	0.15	0.15	0.6	0.6	0.3
Leafy Brassica	0.3	0.15	0.15	0.4	0.4	0.3
Cucurbits	0.3	0.15	0.15	NA	NA	0.3
Eggplant	0.3	0.15	0.15	NA	NA	0.3
Pepper (Bell & Non Bell)	0.3	0.15	0.15	NA	NA	0.3
Tomato	0.3	0.15	0.15	NA	NA	0.3
Head Lettuce	0.3	0.15	0.15	0.6	0.6	0.3
Succulent Peas and Beans	0.3	0.15	0.15	NA	NA	0.3
Dry Peas and Bean	0.3	0.15	0.15	NA	NA	0.3
Turnip Greens	0.3	0.15	0.15	NA	NA	0.3
Root and Tuber	0.3	0.15	0.15	NA	NA	0.3
Treenuts	0.3	0.15	0.15	NA	NA	0.3

Mustang or Fury (1.5 EW or 1.5 EC), Mustang Max (0.8 EC or 0.8 EW), F9043 or any zeta cypermethrin product approved for crop use.

** Any cypermethrin product approved for crop use. NA = Not Applicable.

Maximum Usage When Applying Bifenthrin Products to the Same Crop Within the Same Season

Crop	Maximum Seasonal Total (lbs ai/acre)		
	Bifenthrin		When Applying Bifenthrin Products Plus F9043 to the Same Crop
	F9043	Bifenthrin	
Canola	0.08	0.08	0.08
Field Corn	0.12	0.30	0.30
Sweet Corn	0.18	0.20	0.20
Cotton	0.18	0.50	0.50
Peanut	0.18	0.50	0.50
Potato	0.18	0.50	0.50
Soybean	0.18	0.30	0.30
Head and Stem Brassica	0.18	0.50	0.50
Leafy Brassica	0.18	0.40	0.40
Cucurbits	0.18	0.30	0.30
Eggplant	0.18	0.20	0.20
Pepper (Bell & Non Bell)	0.18	0.20	0.20
Tomato	0.18	0.32	0.32
Head Lettuce	0.18	0.50	0.50
Dry Peas and Beans	0.18	0.30	0.30
Succulent Peas and Beans	0.18	0.20	0.20
Root and Tuber	0.18	0.50 (foliar + soil app.)	0.50
Turnip Greens	0.18	0.40	0.40
Treenuts	0.18	0.50	0.50
Any bifenthrin product approved for crop use			

Field Crops

Canola Crambe and Rapeseed (PHI 35 days)

Pests Controlled	Rate of Application
Cutworm spp Flea beetle	2.5 – 3.5 fl oz/A (0.029 – 0.041 lb ai/A)
Aphid spp Armyworm fall Armyworm southern Armyworm true Armyworm yellowstriped Diamondback moth Flea hopper Grasshopper Looper spp Seedpod weevil Stinkbug spp	3.5 – 4.7 fl oz/A (0.041 – 0.055 lb ai/A)
Restrictions <ul style="list-style-type: none"> • PHI Do not apply within 35 days of harvest • Application Interval Do not make applications less than 14 days apart • Maximum Amount per Application Do not apply more than 4.7 fl oz/A (0.055 lb ai/A) per application • Maximum Amount of F9043 EC Insecticide allowed per Season Do not apply more than 12.8 fl oz/A (0.15 lb ai/A) per season • Refer to the maximum usage tables when applying more than one product containing either zeta cypermethrin or bifenthrin to this crop Remarks Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control. Pyrethroid resistance is common for these pests. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so refer to the resistance management statement in the DIRECTION FOR USE section of this label.	

Field Corn (Grain and Silage) Popcorn Field Corn Grown for Seed
(PHI 30 days for grain and stover and 60 days for forage)
Foliar Use Only

Pests Controlled	Rate of Application
Army cutworm Bean leaf beetle Common stalk borer Cutworm spp Flea beetle spp Green cloverworm Hop vine borer Western bean cutworm	2.5 – 3.5 fl oz/A (0.029 – 0.041 lb ai/A)
Aphid spp Armyworm fall Armyworm southern Armyworm true Armyworm yellowstriped Cereal leaf beetle Chinch bug Corn blotch leafminer (adult) Corn earworm Corn leaf hopper Corn rootworm (adult) Corn silk fly Cucumber beetle (adult) European corn borer False chinch bug Grasshopper spp Greenbug Japanese beetle (adult) Leafhopper spp Meadow spittlebug Sap beetle Southern corn leaf beetle Southwestern corn borer Stinkbug spp Tobacco budworm Webworm spp	3.5 – 4.7 fl oz/A (0.041 – 0.055 lb ai/A)
Restrictions <ul style="list-style-type: none"> • PHI Do not apply within 30 days of harvest for grain and stover and 60 days of harvest for forage • Application Interval Apply at a minimum 3 day intervals or as needed for control • Maximum Amount per Application Do not apply more than 4.7 fl oz/A (0.055 lb ai/A) per application • Maximum Amount of F9043 EC Insecticide allowed per Season Do not apply more than 18.7 fl oz/A (0.22 lb ai/A) per season • Refer to the maximum usage tables when applying more than one product containing either zeta cypermethrin or bifenthrin to this crop • Grazing Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application <p>Use of ultra low volume (ULV) application on corn is prohibited</p> <p>Do not make aerial or ground applications to corn if heavy rainfall is imminent</p>	
Remarks <p>Apply in a minimum of 2.5 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. To improve control by aircraft use 5 gallons of finished spray per acre particularly when initial populations are heavier than normal. When applying by air 1– 2 quarts of emulsified oil may be substituted for 1– 2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.</p> <p>To control ear- attacking pests Apply F9043 EC Insecticide when silking begins and repeat as necessary to maintain control</p>	

Southwestern Corn Borer European Corn Borer Make 2 applications for corn borer control with the first application at or shortly before egg hatch followed by the second application 7 – 10 days later

For control of other insect pests Apply when pests first appear and repeat as necessary

Higher rates will be necessary for heavier initial populations and corn under heat or drought stress

Coverage is essential for control of this pest Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control

Pyrethroid resistance is common for these pests Please consult your local or state agricultural authority to determine if resistance pest populations are in your area If so refer the resistance management statement in the DIRECTION FOR USE section of this label

Sweet Corn (Grain and Silage) Sweet Corn Grown for Seed
(PHI 3 days)
Foliar Use Only

Pests Controlled	Rate of Application
Aphid spp Army cutworm Armyworm fall Armyworm southern Armyworm true Armyworm yellowstriped Cereal leaf beetle Chinch bug Common stalk borer Corn blotch leafminer (adult) Corn earworm Corn rootworm (adult) Corn silk fly Cucumber beetle (adult) Cutworm spp European corn borer False chinch bug Flea beetle Grasshopper spp Greenbug Japanese beetle (adult) Leafhopper spp Sap beetle Southern corn leaf beetle Southwestern corn borer Stinkbug spp Webworm spp Western bean cutworm	3.5 – 4.7 fl oz/A (0.041 – 0.055 lb ai/A)
Restrictions <ul style="list-style-type: none"> • PHI Do not apply within 3 days of harvest • Application Interval Apply at a minimum 3 to 5 day intervals or as needed for control • Maximum Amount per Application Do not apply more than 4.7 fl oz/A (0.055 lb ai/A) per application • Maximum Amount of F9043 EC Insecticide allowed per Season Do not apply more than 28.1 fl oz/A (0.33 lb ai/A) per season • Refer to the maximum usage tables when applying more than one product containing either zeta cypermethrin or bifenthrin to this crop • Grazing Do not graze livestock in treated areas or cut treated crops for feed within 3 days of the last application <p>Use of ultra low volume (ULV) application on corn is prohibited</p> <p>Do not make aerial or ground applications to corn if heavy rainfall is imminent</p> Remarks <p>General Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 20 gallons per acre with ground equipment. When applying by air, 1–2 quarts of emulsified oil may be substituted for 1–2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.</p> <p>To control ear- attacking pests Apply F9043 EC Insecticide when silking begins and repeat as necessary to maintain control. Direct applications at corn ears for optimum control.</p> <p>Southwestern Corn Borer European Corn Borer Make 2 applications for corn borer control with the first application at or shortly before egg hatch followed by the second application 7 – 10 days later.</p> <p>For control of other insect pests Apply when pests first appear and repeat as necessary.</p>	

Higher rates will be necessary for heavier initial populations and corn under heat or drought stress

Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control.

Cotton (PHI 14 days)

Pest Controlled	Rate of Application
Cutworm spp Soybean (banded) thrips Tobacco thrips	2.5 - 3.5 fl oz/A (0.029 - 0.041 lb ai/A)
Armyworm fall Armyworm yellowstriped Boll weevil Bollworm Cabbage looper Cotton aphid Cotton fleahopper Cotton leafperforator European corn borer Grasshopper spp Saltmarsh caterpillar Southern garden leafhopper Stinkbug spp Tobacco budworm	3.5 - 4.7 fl oz/A (0.041 - 0.055 lb ai/A)
Restrictions <ul style="list-style-type: none"> • PHI Do not apply within 14 days of harvest • Maximum Amount per Application Do not apply more than 4.7 fl oz/A (0.055 lb ai/A) per application • Maximum Amount of F9043 EC Insecticide allowed per Season Do not apply more than 28.1 fl oz/A (0.33 lb ai/A) per season • Refer to the maximum usage tables when applying more than one product containing either zeta cypermethrin or bifenthrin to this crop • Grazing Do not graze livestock in treated areas or cut treated crops for feed 	
Remarks F9043 EC Insecticide may be applied in water or refined vegetable oil (soybean/cottonseed)	
Application in Water Apply in a minimum of 5 gallons per acre with ground equipment or 1 gallon per acre by aircraft. When applying by air, 1 quart of emulsified oil may be substituted for one quart of water in the finished spray.	
ULV Application Apply the recommended rate of F9043 EC Insecticide in refined vegetable oil in a minimum of 1 quart of finished spray per acre with aircraft calibrated to give adequate coverage.	
To Control Boll Weevil Apply F9043 EC Insecticide at an interval of 3 to 4 days until pest numbers are reduced to acceptable levels.	
To Control Aphids Apply when pests first appear. Repeat as necessary to maintain control.	
Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control.	
Pyrethroid resistance is common for these pests. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so, refer the resistance management statement in the DIRECTION FOR USE section of this label.	

Peanut (PHI 14 days)

Pests Controlled	Rate of Application
Aphid spp Armyworm fall Armyworm southern Armyworm true Armyworm yellowstriped Bean leaf beetle Corn earworm Cutworm spp Grasshopper spp Green cloverworm Leafhopper spp Lesser cornstalk borer Looper spp Red necked peanut worm Southern corn rootworm (adult) Stinkbug spp Threecornered alfalfa hopper Vegetable weevil Velvetbean caterpillar Whitefringed beetle (adult)	3.5 – 4.7 fl oz/A (0.041 – 0.055 lb ai/A)
Restrictions <ul style="list-style-type: none"> • PHI Do not apply within 14 days of harvest • Application Interval Do not make applications less than 14 days apart • Maximum Amount per Application Do not apply more than 4.7 fl oz/A (0.055 lb ai/A) per application • Maximum Amount of F9043 EC Insecticide allowed per Season Do not apply more than 28.1 fl oz/A (0.33 lb ai/A) per season Restricted to 6 foliar applications per season • Refer to the maximum usage tables when applying more than one product containing either zeta cypermethrin or bifenthrin to this crop Remarks Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). Follow appropriate spray drift precautions on this label. Do not feed green immature plants and peanut hay to livestock. Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control.	

Potato (PHI 21 days)

Pests Controlled	Rate of Application
Cutworm spp Flea beetle spp	2.5 – 3.5 fl oz/A (0.041 – 0.029 lb ai/A)
Aphid spp Armyworm fall Armyworm southern Armyworm true Armyworm yellowstriped Banded cucumber beetle Chinch bug Colorado potato beetle Cucumber beetle (adult) European cornborer False chinch bug Grasshopper spp Looper spp Potato leafhopper Sugarcane beetle Sweetpotato flea beetle Sweetpotato weevil (adult) Potato tuberworm	3.5 – 4.7 fl oz/A (0.041 – 0.055 lb ai/A)
Restrictions <ul style="list-style-type: none"> • PHI Do not apply within 21 days of harvest • Application Interval Do not make applications less than 21 days apart • Maximum Amount per Application Do not apply more than 4.7 fl oz/A (0.055 lb ai/A) per application • Maximum Amount of F9043 EC Insecticide allowed per Season Do not apply more than 28.1 fl oz/A (0.33 lb ai/A) per season Restricted to 2 foliar applications per season • Refer to the maximum usage tables when applying more than one product containing either zeta cypermethrin or bifenthrin to this crop <p>Leaves cannot be used for food or feed</p> Remarks Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 3 gallons by air). Follow appropriate spray drift precautions on this label. Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control. For Tuberworm control (adults and larvae) apply prior to harvest or senescence when economic thresholds are met. Only tuberworms that have not bored or tunneled into the plant can be controlled with a foliar application of F9043 EC Insecticide.	

Soybeans (PHI 21 days)

Pest Controlled	Rate of Application
Cutworm spp Flea beetle spp Green cloverworm Painted lady (thistle) caterpillar Silverspotted skipper	2.5 - 3.5 fl oz/A (0.029 - 0.041 lb ai/A)
Alfalfa caterpillar Armyworm fall Armyworm southern Armyworm true Armyworm yellowstriped Bean leaf beetle Blister beetle spp Corn earworm Corn rootworm (adult) Cowpea curculio Cucumber beetle (adult) Dectes stem borer (adult) European corn borer False chinch bug Grape colaspis (adult) Grasshopper spp Hornworm spp Imported cabbageworm Japanese beetle (adult) Leaf skeletonizer spp Leafhopper spp Leafminer spp (adult) Lesser cornstalk borer Looper spp Mexican bean beetle Pea leaf weevil Saltmarsh caterpillar Seedcorn maggot (adult) Soybean aphid Spittlebug Stinkbug spp Threecornered alfalfa hopper Thrips Tobacco budworm Velvetbean caterpillar Webworm	3.5 - 4.7 fl oz/A (0.041 - 0.055 lb ai/A)
Restrictions <ul style="list-style-type: none"> • PHI Do not apply within 21 days of harvest • Application Interval Do not make applications less than 30 days apart • Maximum Amount per Application Do not apply more than 4.7 fl oz/A (0.055 lb ai/A) per application • Maximum Amount of F9043 EC Insecticide allowed per Season Do not apply more than 28.1 fl oz/A (0.33 lb ai/A) per season • Refer to the maximum usage tables when applying more than one product containing either zeta cypermethrin or bifenthrin to this crop • Grazing Do not graze or harvest treated soybean forage straw or hay for livestock feed Remarks Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). Follow appropriate spray drift precautions on this label. Coverage is essential for control of this pest. Under heavy outbreak conditions, tank mixing with another product that is labeled for this pest is recommended for control.	

* For optimum control apply F9043 EC Insecticide 10 days after adult emergence. Follow 10-14 days later with an application of a product containing zeta cypermethrin as the sole active ingredient such as Mustang Max Insecticide applied at the single maximum application rate under the soybean use directions for that product.

Pyrethroid resistance is common for this pest. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so, refer the resistance management statement in the *DIRECTION FOR USE* section of this label.

Vegetable and Small Fruit Crops

Brassica Vegetables

Head & Stem Brassica (PHI 7 days)

Including Broccoli broccoli Chinese Brussels sprouts cabbage
cabbage Chinese (napa) cabbage Chinese mustard cauliflower
cavalo broccolo kohlrabi

Pest Controlled	Rate of Application
Aphid spp Armyworm fall Armyworm southern Armyworm true Armyworm yellowstriped Click beetle (wireworm adult) Corn earworm Crickets Cucumber beetle Cutworm spp Diamondback moth Flea beetle spp Grasshopper spp Imported cabbageworm Leafhopper spp (adult) Leafminer spp Looper spp Saltmarsh caterpillar Southern cabbageworm Stinkbug spp Tobacco budworm	3.5 – 4.7 fl oz/A (0.041 – 0.055 lb ai/A)
Restrictions <ul style="list-style-type: none"> • PHI Do not apply within 7 days of harvest • Application Interval Do not make applications less than 7 days apart • Maximum Amount per Application Do not apply more than 4.7 fl oz/A (0.055 lb ai/A) per application • Maximum Amount of F9043 EC Insecticide allowed per Season Do not apply more than 28.1 fl oz/A (0.33 lb ai/A) per season. Do not make more than 5 applications after bloom • Refer to the maximum usage tables when applying more than one product containing either zeta cypermethrin or bifenthrin to this crop Remarks Apply in a minimum of 5 gallons of finished spray per acre by air or in a minimum of 15 gallons per acre with ground equipment. When applying by air, 1–2 quarts of emulsified oil may be substituted for 1–2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Coverage is essential for control of this pest. Under heavy outbreaks, tank mixing with another product that is labeled for this pest is recommended for control. Pyrethroid resistance is common for this pest. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so, refer the resistance management statement in the DIRECTION FOR USE section of this label.	

Leafy Brassica (PHI 7 days)

Including Broccoli raab cabbage Chinese (bok choy) collards kale mizuna mustard greens mustard spinach rape greens

Pest Controlled	Rate of Application
Aphid spp Armyworm fall Armyworm southern Armyworm true Armyworm yellowstriped Cabbageworm Click beetle (wireworm adult) Corn earworm Crickets Cucumber beetle (adult) Cutworm spp Diamondback moth Flea beetle spp Grasshopper spp Imported cabbageworm Japanese beetle (adult) Leafhopper spp Leafminer (adult) Looper spp Saltmarsh caterpillar Stinkbug spp Onion thrips Tobacco budworm Wireworm (adult)	3.5 – 4.7 fl oz/A (0.041 – 0.055 lb ai/A)
Restrictions • PHI Do not apply within 7 days of harvest • Application Interval Do not make applications less than 7 days apart • Maximum Amount per Application Do not apply more than 4.7 fl oz/A (0.055 lb ai/A) per application • Maximum Amount of F9043 EC Insecticide allowed per Season Do not apply more than 28.1 fl oz/A (0.33 lb ai/A) per season • Refer to the maximum usage tables when applying more than one product containing either zeta cypermethrin or bifenthrin to this crop Remarks Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 15 gallons by ground and 5 gallons by air). Follow appropriate spray drift precautions on this label. Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product which is labeled for this pest is recommended for control. Pyrethroid resistance is common for this pest. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so refer the resistance management statement in the DIRECTION FOR USE section of this label.	